

IUE OBSERVATIONS OF THE SOCKET STAR MT Ori AND ITS CIRCUMSTELLAR ENVIRONMENT

BRUCE MCCOLLUM
Astronomy Programs, Computer Sciences Corporation

and

MICHAEL CASTELAZ
East Tennessee State University

7N-89
53157
P-3

Abstract. We have obtained a series of large aperture images taken at different spatial locations relative to MT Ori which form a "cross-cut" of about 50" in length through the star and its immediate environment. The data show a marked drop in nebular continuum emission and 1909 Å line emission, centered on MT Ori, which is consistent with the appearance of MT Ori on high-quality optical plates as being embedded in a region of low H-alpha brightness, which may be physically associated with MT Ori.

Key words: socket stars-circumstellar material-pre-main sequence stars.

1. Introduction

Socket stars are stars which appear on high-quality plates of HII regions to be embedded in dark regions 5" to 20" in radius, approximately centered on the star, called "sockets" (Glushkov 1952, 1986, 1991; Feibelman 1989). Feibelman (1989) reports that "dozens" of sockets are visible in Orion and other HII regions. Although some are likely to be chance superpositions of stars against physically unrelated dark background regions, Feibelman (1989) argues that overall socket stars occur more frequently than can be accounted for by chance superpositions. Almost all Orion socket stars show an IR excess (Castelaz 1990), although this is common for non-socket stars in the Orion region. It is reasonable to suspect that the dark sockets represent a circumstellar envelope of some sort, possibly the remnant of the "cocoon" in which the star was formed. In an effort to obtain more information as to the physical nature of the sockets, we undertook IUE observations of one socket star in Orion.

2. Program and Results

We obtained a series of large aperture SWP images of the MT Ori vicinity. Each image was taken with the telescope pointed in a slightly different direction, so that the images overlapped along the short axis of the large aperture (10" x 20" oval). The sequence of images forms a cross-cut in the NW to SE direction. A few images were also taken NE of MT Ori. Because MT Ori is 10th magnitude, and has spectral type F with an $E(B-V) = 2.0$ (Glushkov, 1993, private communication), the detected flux in our spectra (which are short exposures) must be from nebular

W. Wunsteler et al. (eds.), *Frontiers of Space and Ground-Based Astronomy*, 561-562.

(NASA-CR-189401) IUE OBSERVATIONS
OF THE SOCKET STAR MT ORI AND ITS
CIRCUMSTELLAR ENVIRONMENT
(Computer Sciences Corp.) 3 p

N95-71365

Unclass




Fig. 1. Observed continuum and 1909 Å CIII] line flux changes in SE (left) to NW (right) direction across MT Ori. Continuum bins were 50 Å wide.

emission rather than MT Ori. We note that MT Ori is thought to be a pre-main sequence star on the basis of its X-ray emission (Vaiana et al. 1981).

Despite the limited spatial resolution imposed by the dimensions of the large aperture, we find a clear local minimum in nebular emission which appears to be centered on MT Ori (Fig. 1). This area of diminished emission has a radius of about 20". We do not know the UV emission SW of MT Ori, but the available fluxes do possess spatial symmetry on three sides of MT Ori, suggesting a "socket" in the UV. Although the data do not rule out the possibility that this is the result of a chance superposition of MT Ori against a relatively faint area of the nebula which is physically unrelated to the star, the fact that the UV emission reaches its minimum coincident with the position of MT Ori and the fact that the flux changes monotonically on either side suggests that the star is physically related to the local nebular minimum.

References

- Castelaz, M.W. 1990, *Ap.J.*, 348, L29
 Feibelman, W. 1989, *PASP*, 101, 547
 Glushkov, Yu.I. 1952, *Trudy Astrophys. Inst. Acad. Kazakh SSR*, No. 1449
 Glushkov, Yu.I. 1986, *Astron. Tsirk.*, No. 1449
 Glushkov, Yu.I. 1991, *Astron. and Astrophysical Transactions*, 1, 75
 Vaiana, G.S., Cassinelli, J.P., Fabbiano, G., Giacconi, R., Golub, L., Gorenstein, P., Haisch, B.M., Harnden, F.R., Johnson, H.M., Linsky, J.L., Maxson, C.W., Mewe, R., Rosner, R., Seward, F., Topka, K., & Zwaan, C. 1981, *ApJ*, 245, 163

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE November 1994	3. REPORT TYPE AND DATES COVERED Contractor Report	
4. TITLE AND SUBTITLE IUE Observations of the Socket Star MT Ori and It's Circumstellar Environment			5. FUNDING NUMBERS Code 684.1 7N-89 53157	
6. AUTHOR(S) Bruce McCollum (CSC) and Michael Castelaz (East Tennessee State University)				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Computer Sciences Corporation System Science Division 4061 Powder Mill Road Calverton, MD 20705			8. PERFORMING ORGANIZATION REPORT NUMBER S-97223-E	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) NASA Aeronautics and Space Administration Washington, D.C. 20546-0001			10. SPONSORING/MONITORING AGENCY REPORT NUMBER CR-189401	
11. SUPPLEMENTARY NOTES Technical Monitor: D. West, Code 684.1				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Unclassified-Unlimited Subject Category: 89 Report available from the NASA Center for AeroSpace Information, 800 Elkridge Landing Road, Linthicum Heights, MD 21090; (301) 621-0390.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) We have obtained a series of large aperture images taken at different spatial locations relative to MT Ori which form a "cross-cut" of about 50" in length through the star and its immediate environment. The data show a marked drop in nebular continuum emission and 1909 Å line emission centered on MT Ori, which is consistent with the appearance of MT Ori on high-quality optical plates as being embedded in a region of low H-alpha brightness, which may be physically associated with MT Ori.				
14. SUBJECT TERMS Socket stars, Circumstellar material, Pre-main sequence stars			15. NUMBER OF PAGES 2	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	